

## II.4 Existence of covering spaces

### 1. Existence of a covering space assuming the existence of a universal covering space.


$$\begin{array}{c}
 (\tilde{X}, \tilde{x}) \\
 \downarrow p : \text{a universal covering with a deck transformation group } G \xrightarrow{\theta(\cong)} \pi_1(X, x) \\
 (X, x)
 \end{array}$$

정리 1 If  $H < G$ , then

$$\begin{array}{ccc}
 (\tilde{X}, \tilde{x}) & & \\
 & \searrow q & \\
 p \downarrow & & (\tilde{X}_H, \tilde{x}) \\
 & \swarrow r & \\
 (X, x) & & 
 \end{array}$$

where  $\tilde{X}_H = H \backslash \tilde{X}$  with  $r_{\#} \pi_1(\tilde{X}_H, \tilde{x}) = \theta^{-1}(H)$  and the quotient map  $q$  and the induced map  $r$  become covering maps.

증명



왼쪽 그림에서와 같이  $p$ 에 의해 evenly cover되는  $U$ 를 잡으면  $H$ 는  $p^{-1}(U)$ 에 permutation으로 작용하므로  $H$  action에 대한 quotient map  $q$ 가 covering map이 되는 것은 분명하고  $r$ 을  $q$ 에 의해  $p$ 로부터 induced되는  $\tilde{p}$ 로 정의

하면  $r$ 은 당연히 onto이고,  $U$ 가 역시  $r$ 에 의해서 evenly covered 되므로  $r$ 은 covering 이 된다. 그리고

$$\begin{aligned} \alpha \in \theta^{-1}(H) \subset \pi_1(X, x) &\Leftrightarrow \theta(\alpha) \in H \Leftrightarrow \tilde{x} \cdot \alpha \in H \cdot \tilde{x} \\ \Leftrightarrow \bar{x} := f(\tilde{x}) = q(\tilde{x} \cdot \alpha) = \bar{x} \cdot \alpha &\Leftrightarrow \alpha \in \Pi_{\bar{x}} = r_{\sharp} \pi_1(\tilde{X}_H, \bar{x}) \text{ 가 성립하므로,} \\ r_{\sharp} \pi_1(\tilde{X}_H, \bar{x}) &= \theta^{-1}(H) \text{ 이다.} \end{aligned}$$

□

## 2. Existence of a universal covering space.

Idea : the set of homotopy classes of paths from  $x$  to  $y$   
 $\xleftrightarrow[1-1]{p^{-1}(y) \subset \tilde{X}}$ , universal covering, via  $[\alpha] \leftrightarrow \tilde{\alpha}(1)$

**정의 1** A space  $X$  is said to be **semilocally simply connected** if for each  $x \in X$ , there is a neighborhood  $U$  of  $x$  such that the homomorphism

$$i_{\sharp} : \pi_1(U, x) \rightarrow \pi_1(X, x)$$

induced by inclusion is trivial.

Assume that  $X$  is path-connected, locally path-connected and semilocally simply connected.

Let  $(P, x) = \{\alpha : I \rightarrow X \mid \alpha(0) = x\}$

Define  $\tilde{X} = (P, x) / \sim$  (recall  $\alpha \sim \beta \Leftrightarrow \alpha \simeq \beta \text{ rel } \partial$ ) and  $p : \tilde{X} \rightarrow X$  by  $p([\alpha]) = \alpha(1)$

### Topology of $\tilde{X}$

For  $\alpha \in (P, x)$  and open set  $U$  with  $\alpha(1) \in U$ , let  $(\alpha, U) := \{\alpha * \alpha' \mid \alpha' : I \rightarrow U \text{ with } \alpha'(1) = \alpha(0)\} / \sim \subset \tilde{X}$ .

이때 주어진  $y = \alpha(1) \in X$ 에 대해,  $U$ 를 path-connected, semilocally simply connected neighborhood of  $y (= \alpha(1))$  라고 하면  $p| : (\alpha, U) \rightarrow U$ 는 onto 이고 one-to-one 이다.

이제  $p$ 가 evenly covered 임을 보이자. 즉,  $p^{-1}(U) = \coprod_{\alpha(1)=y} (\alpha, U)$ :

$(\alpha, U) \cap (\beta, U) \neq \emptyset$  라고 하자.

$\gamma \in (\alpha, U) \cap (\beta, U)$ 에 대하여  $\alpha'$ 와  $\beta'$ 가 존재해서

$$\begin{aligned} \alpha * \alpha' \sim \gamma \sim \beta * \beta' \\ \Rightarrow \alpha * \alpha' * \bar{\alpha}' \sim \beta * \alpha' * \bar{\alpha}' (\because U \text{가 semilocally simply connected 이므로}) \\ \beta * \beta' \sim \beta * \alpha' \end{aligned}$$

$$\Rightarrow \alpha \sim \beta$$

$$\therefore (\alpha, U) \cap (\beta, U) \neq \emptyset \quad \Rightarrow \quad (\alpha, U) = (\beta, U)$$

Take  $\{(\alpha, U) | U : \text{open neighborhood of } \alpha(1), \alpha \in (P, x)\}$  as a base for a topology of  $\tilde{X}$ .

**Check**

1.  $\bigcup(\alpha, U) = \tilde{X}$  (obvious)

2.  $\gamma \in (\alpha, U) \cap (\beta, U)$

$\Rightarrow \exists W \subset U \cap V$  such that  $(\gamma, W) \subset (\alpha, U) \cap (\beta, U)$  여기서  $W$ 는 path-connected neighborhood of  $\gamma(1)$ 이고  $U$ 가 semilocally simply connected 이므로 subset  $W$ 도 마찬가지로 (obvious)

$\tilde{X}$ 가 path-connected 임을 보이자.

임의의  $[\alpha] \in \tilde{X}$ 에 대해서  $\alpha_s(t) := \alpha(st)$ 라고 하면  $\alpha_s$ 는  $\alpha$ 와  $x$ (constant path)를 잇는 path이다. 여기서  $\tilde{\alpha}(s) := [\alpha_s]$ 라고 정의하면  $\tilde{\alpha}(0) = [x], \tilde{\alpha}(1) = [\alpha]$ 가 되어  $[\alpha]$ 와  $[x]$ 는 path 로 연결된다. (exercise :  $\tilde{\alpha}$  is continuous)

마지막으로  $\tilde{X}$ 가 simply connected 임을 보이자.

Let  $\tau$  be a loop in  $(\tilde{X}, \tilde{x})$ , where  $\tilde{x} = [x]$

$$\Rightarrow \alpha := p \circ \tau \text{ is a loop in } X \text{ and } \tilde{\alpha} = \tau$$

$$\Rightarrow [\alpha] = \tilde{\alpha}(1) = \tau(1) = [x] \quad \Rightarrow \quad \alpha \sim x \quad \Rightarrow \quad \tau \sim \tilde{x} \quad \blacksquare$$

**따름정리 2**  $\forall H < \pi_1(X, x), \exists$  a covering space  $(\tilde{X}, \tilde{x})$  corresponding to  $H$ , i.e.  $p_{\#}\pi_1(\tilde{X}, \tilde{x}) = H$

**Remark**

1. Universal covering is "universal", i.e. it covers every other covering by lifting theorem and universal covering is clearly unique up to isomorphism.

2.  $X$  has a universal covering.

$\Rightarrow X$  is semilocally simply connected.

$$\begin{array}{ccc} \pi_1(\tilde{U}, \tilde{x}) & \xrightarrow{i_{\#}} & \pi_1(\tilde{X}, \tilde{x}) = 0 \\ \cong \downarrow p_{\#} & \circlearrowleft & \downarrow p_{\#} \\ \pi_1(U, x) & \xrightarrow{i_{\#}} & \pi_1(X, x) \end{array}$$

로 부터 당연하다.

**숙제 8**

Let  $(G, e)$  be a topological group and  $p : (\tilde{G}, \tilde{e}) \rightarrow (G, e)$  be a covering. Then we can lift the group structure of  $G$  to  $\tilde{G}$  so that  $p$  becomes a homomorphism unique up to the choice of identity  $\tilde{e} \in p^{-1}(e)$ .